

Targeting MRGPRX2 with venom peptide P17: A therapeutic avenue for DSS-Induced colitis

Karthi Duraisamy,^{1,2} Mukesh Kumar,¹ Abdullah Nawabjan,¹ Emily Kwun Kwan Lo,¹ Benjamin Lefranc,² Elsa Bonnafé,³ Michel Treillou,³ Hani El-Nezami,^{1,4} Billy K.C. Chow,^{*1} Jérôme Leprince.^{*2}


¹School of Biological Sciences, The University of Hong Kong, Hong Kong, China; ²INSERM U1239 NordIC, PRIMACEN, Université Rouen Normandie, Rouen, France; ³EA7417 B7SB, Université Fédérale Toulouse Midi-Pyrénées, INU Champollion, Albi, France; ⁴Institute of Public Health and Clinical Nutrition, University of Eastern Finland, Kuopio, Finland.

Karthi.duraisamy@univ-rouen.fr


<https://doi.org/10.17952/37EPS.2024.P2154>

Introduction

Inflammatory Bowel Disease



Common Symptoms of IBD





Peptide P17

Novel Anti Microbial Peptide (AMP) from the venom of *Tetramorium bicarinatum*


LFKEILEKIKAKL-NH2

Known to activate an unknown GPCR






Results & discussion


Identification of P17 receptor




Group	GPCRs
Class A	262
Class B	16
Class C	17
Adhesion	17
Other	2
Total	314



Structure-Activity Relationship of P17 and MRGPRX2 Activation




MRGPRX2 Activation in Mast Cells Promotes Monocyte Recruitment and Differentiation




Acknowledgements & affiliations




MRGPRX2/B2 deficiency exacerbates colitis in DSS mouse model




MRGPRB2 ameliorates severity of colitis in DSS mouse model



MRGPRB2 ligands reduce inflammatory cytokines and immune cell infiltration in DSS-Induced Colitis mice model



Conclusion & future prospective



References

- Duraisamy, K., Singh, K., Kumar, M., Lefranc, B., Bonnafé, E., Treillou, M., ... & Chow, B. K. (2022). P17 induces chemotaxis and differentiation of monocytes via MRGPRX2-mediated mast cell activation. *Journal of Allergy and Clinical Immunology*, 149(11), 275-291.
- Duraisamy, K., Kumar, M., Nawabjan, A., Lo, E. K. K., Hui, L. H., Lefranc, B., ... & Chow, B. K. (2024). MRGPRB2/X2 and the analogous effects of its agonist and antagonist in DSS-induced colitis in mice. *Biomedicine & Pharmacotherapy*, 174, 116471.
- Kumar, M., Duraisamy, K., Anandapany, R. R., Chan, C. S., & Chow, B. K. (2023). Novel small molecule MRGPRX2 antagonists inhibit a murine model of allergic reaction. *Journal of Allergy and Clinical Immunology*, 151(4), 1110-1122.
- do Nascimento, R. D. P., da Fonseca Machado, A. P., Galvez, J., Caziani, C. B. B., & Junior, M. R. M. (2020). Ulcerative colitis: Gut microbiota, immunopathogenesis and application of natural products in animal models. *Life sciences*, 248, 118129.
- Lakhan, S. E., & Kirchgessner, A. (2010). Neuroinflammation in inflammatory bowel disease. *Journal of neuroinflammation*, 7, 1-12.
- Tolley, S., Valaigga, R., Anderson, L., Cannon, A. R., Choudhry, M. A., & Campbell, E. M. (2021). DSS-induced inflammation in the colon drives a proinflammatory signature in the brain that is ameliorated by prophylactic treatment with the S100A9 inhibitor poaquimod. *Journal of Neuroinflammation*, 18, 1-14.
- Benmoussa, K., Authier, H., Prat, M., Alédidine, M., Lefèvre, L., Rahabi, M., C., ... & Coste, A. (2017). P17, an original host defense peptide from an ant venom, promotes anti-fungal activities of macrophages through the induction of C-type lectin receptors dependent on L184-mediated PF4γ activation. *Frontiers in Immunology*, 8, 1450.